This is one unit... For ma

For more detailed information on calculating units see - www.nhs.uk/Livewell/alcohol/Pages/alcohol-units.aspx



Adults who don't regularly Increased relaxation AUDIT-C* Low risk 0 - 4drinking drink more than 14 units per Sociability ASSIST-Lite* 0-4 week, spread over three or · Sensory enjoyment of AUDIT *** 0-7 alcoholic drinks more days, are at low risk of harm from drinking. Drinking in this way raises AUDIT-C* 5-7 Increasing Progressively your long-term risk of ill ASSIST-Lite** 5-7 risk increasing risk of: AUDIT *** health. 8-15 drinking Low energy Relationship problems Drinking in this way is likely AUDIT-C* Higher 8-10 Depression ASSIST-Lite** 8-10 to be affecting your health risk Insomnia AUDIT *** or wellbeing in some way 16-19 drinking Impotence already. Injury AUDIT-C* Your drinking may have 11-12 Possibly · High blood pressure ASSIST-Lite** 11-12 become quite problematic dependent · Breast, mouth and AUDIT *** and further assessment by 20 - 40throat cancers an alcohol specialist is Alcohol dependence advised. Liver disease

*Short 3 question AUDIT questionnaire. **ASSIST-Lite seven-item drug, alcohol and tobacco screening questionnaire, ***Full 10 question AUDIT questionnaire.

NHS

There is no completely safe level of drinking and drinking even small amounts of alcohol can incur risk in certain circumstances

For example, with strenuous exercise, operating heavy machinery, driving or if you are on certain medications.

If you are pregnant or planning a pregnancy, the safest approach is not to drink alcohol at all.


Drinking in pregnancy can harm the baby, with the more you drink the greater the risk.

The risk of harm to the baby is likely to be low if a woman has drunk only small amounts of alcohol before she knew she was pregnant or during pregnancy.

This leaflet is based on the "**How Much Is Too Much?"** Simple Structured Advice Intervention Tool, developed by Newcastle University and the Drink Less materials originally developed at the University of Sydney as part of a W.H.O. collaborative study.

Download this alcohol advice tool from https://app.box.com/v/CQUINstructured-advice-tool

What's everyone else like?



The potential benefits of cutting down

Psychological/Social/Financial

- Improved mood
- Improved relationships
- More time for hobbies and interests
- Reduced risk of drink driving
- Save money

Physical

- Sleep betterMore energy
- More energy
- Lose weight
- Reduced risk of injury
- Improved memory
- Better physical shape
- Reduced risk of high blood pressure
- Reduced risk of cancer
- Reduced risk of liver disease

Reduced risk of brain damage

Making your plan

- Have several 'drink-free' days, when you don't drink at all
- When you do drink, set yourself a limit and stick to it
- Quench your thirst with non-alcohol drinks before and in-between alcoholic drinks
- Avoid drinking in rounds or in large groups
- Eat when you drink have your first drink after starting to eat
- Switch to lower alcohol beer/lager
- Avoid going to the pub after work
- Plan activities and tasks at those times you would usually drink
- When bored or stressed do something physical instead of drinking
- Avoid or limit the time spent with "heavy" drinking friends

What targets should you aim for?

There is no completely safe level of drinking, but by sticking within these guidelines, you can lower your risk of harming your health:

- Adults are advised not to regularly drink more than 14 units a week
- If you do drink as much as 14 units in a week, spread this out evenly over 3 or more days.

What's your personal target?



Some brief advice about alcohol and your health



DRINK FREE DAYS APP

Choose your Drink Free Days and get reminders, support and practical advice to change your drinking habits for good.

DOWNLOAD THE APP www.nhs.uk/oneyou/apps

More information is available from One You: www.nhs.uk/oneyou