

In Oldham 199 5-year-olds (approximately 60.5% of those sampled) were examined at school by trained and calibrated examiners using the national standard method².

Figure 1: Prevalence of experience of dental decay and mean number of teeth with experience of dental decay in 5-year-olds in Oldham, other local authorities in the North West and England.




Table 1: Experience of dental decay in 5-year-olds in Oldham, other local authorities in the North West and England.

| Local authority | Prevalence of experience of dental decay (%) | Mean number of teeth with experience of dental decay in all examined children n (95% confidence intervals) | Mean number of teeth with experience of dental decay in children with any decay experience n (95% confidence intervals) |
|-----------------------|--|--|---|
| Blackburn with Darwen | 50.9 | 2.3 (1.87 - 2.82) | 4.6 (3.94 - 5.29) |
| Oldham | 43.2 | 1.9 (1.45 - 2.34) | 4.4 (3.63 - 5.16) |
| Rochdale | 40.7 | 1.7 (1.30 - 2.19) | 4.3 (3.64 - 4.93) |
| Salford | 39.0 | 1.6 (1.25 - 2.01) | 4.2 (3.48 - 4.88) |
| Manchester | 38.3 | 1.8 (1.43 - 2.10) | 4.6 (3.98 - 5.24) |
| Liverpool | 37.0 | 1.4 (1.05 - 1.69) | 3.7 (3.09 - 4.33) |
| Knowsley | 35.4 | 1.3 (0.93 - 1.75) | 3.8 (2.94 - 4.63) |
| Bury | 35.2 | 1.4 (1.07 - 1.66) | 3.9 (3.35 - 4.44) |
| Blackpool | 33.4 | 1.1 (0.74 - 1.56) | 3.4 (2.50 - 4.35) |
| Tameside | 33.1 | 1.0 (0.75 - 1.31) | 3.1 (2.52 - 3.71) |
| Bolton | 32.7 | 1.1 (0.84 - 1.45) | 3.5 (2.83 - 4.18) |
| Wigan | 31.9 | 1.0 (0.79 - 1.26) | 3.2 (2.70 - 3.74) |
| NORTH WEST | 31.7 | 1.2 (1.14 - 1.25) | 3.8 (3.65 - 3.89) |
| Lancashire | 30.4 | 1.2 (1.06 - 1.25) | 3.8 (3.59 - 4.02) |
| Sefton | 29.4 | 1.0 (0.67 - 1.30) | 3.4 (2.61 - 4.10) |
| Halton | 27.0 | 1.0 (0.72 - 1.36) | 3.8 (3.01 - 4.67) |

| Local authority | Prevalence of experience of dental decay (%) | Mean number of teeth with experience of dental decay in all examined children n (95% confidence intervals) | Mean number of teeth with experience of dental decay in children with any decay experience n (95% confidence intervals) | |
|---------------------------|--|---|---|--|
| Trafford | 26.0 | 1.0 (0.72 - 1.32) | 3.9 (3.05 - 4.77) | |
| Warrington | 24.3 | 0.8 (0.52 - 1.01) | 3.1 (2.43 - 3.85) | |
| Cumbria | 24.2 | 0.8 (0.67 - 0.86) | 3.2 (2.89 - 3.43) | |
| ENGLAND | 23.4 | 0.8 (0.78 - 0.81) | 3.4 (3.36 - 3.44) | |
| Cheshire West and Chester | 22.7 | 0.6 (0.41 - 0.78) | 2.6 (2.07 - 3.19) | |
| Stockport | 22.0 | 0.7 (0.43 - 0.94) | 3.1 (2.21 - 3.99) | |
| Cheshire East | Did Not Participate | | | |
| St. Helens | Did Not Participate | | | |
| Wirral | Did Not Participate | | | |

Table 2. Measures of oral health among 5-year-olds in Oldham, it's statistical neighbours, i the North West and England.

| | Oldham | Statistical neighbour within North West: Rochdale | Statistical neighbour comparator 1: Rochdale | North West | England |
|--|--------|--|---|---------------|---------|
| Prevalence of experience of dental decay | 43.2% | 40.7% | 40.7% | 31.7% | 23.4% |
| Mean number of teeth with experience of dental decay | 1.9 | 1.7 | 1.7 | 1.2 | 0.8 |
| Mean number of teeth with experience of decay in those with experience of dental decay | 4.4 | 4.3 | 4.3 | 3.8 | 3.4 |
| Mean number of decayed teeth in those with experience of dental decay | 3.8 | 3.6 | 3.6 | 3.0 | 2.7 |
| Proportion with active decay | 41.0% | 37.6% | 37.6% | 29.0% | 20.4% |
| Proportion with experience of tooth extraction ⁱⁱ | 4.5% | 3.0% | 3.0% | 2.9% | 2.2% |
| Proportion with dental abscess | 1.7% | 1.8% | 1.8% | 1.2% | 1.0% |
| Proportion with teeth decayed into pulp | 12.3% | 9.2% | 9.2% | 5.5% | 3.3% |
| Proportion with decay affecting incisorsiii | 12.6% | 11.4% | 11.4% | 7.7% | 5.2% |
| Proportion with high levels of plaque present on upper front teeth ^{iv} | 1.7% | 5.2% | 5.2% | 2.2% | 1.2% |


¹ generated by the children's services statistical neighbour benchmarking tool, the neighbour within the North West has "Very Close" comparator characteristics and the national neighbour 1 has "Very Close" comparator characteristics³.

ii experience of extraction of one or more teeth on one or more occasions.

iii dental decay involving one or more surfaces of upper anterior teeth.


iv indicative of poor tooth brushing habits.

Figure 2: Prevalence of experience of dental decay in 5-year-olds in Oldham, by local authority Index of Multiple Deprivation (IMD) 2019 quintiles.




Error bars represent 95% confidence limits

Figure 3: Slope index of inequality in the prevalence of experience of dental decay in 5-year-olds in the North West.



Error bars represent 95% confidence limits

Figure 4: Prevalence of experience of dental decay in 5-year-olds in Oldham, the North West and England, by year.



Error bars represent 95% confidence limits

Table 3: Experience of dental decay in 5-year-olds in the North West, by ethnic group.

| Ethnic group | Number of children examined (N) | Prevalence of experience of dental decay (%) | Mean number of teeth with experience of dental decay among children with any experience of dental decay n (95% CI) | Prevalence of dental decay affecting incisors (%) |
|--------------------------|--|---|---|--|
| Other ethnic background | 78 | 67.9 | 5.0 (4.05 - 5.98) | 30.8 |
| Asian / Asian British | 803 | 50.4 | 4.7 (4.39 - 5.07) | 20.0 |
| Not provided | 97 | 41.2 | 3.8 (2.59 - 4.96) | 9.3 |
| Black / Black British | 200 | 38.5 | 3.9 (3.25 - 4.62) | 12.5 |
| Mixed | 223 | 32.3 | 3.4 (2.74 - 4.10) | 9.0 |
| White | 6,753 | 26.7 | 3.4 (3.25 - 3.51) | 4.9 |
| North West | 8,154 | 31.7 | 3.8 (3.65 - 3.89) | 7.7 |

Figure 5: Prevalence of experience of dental decay in 5-year-olds in the North West, by ethnic group.




Table 4: Prevalence and severity of experience of dental decay in 5-year-olds in Oldham, by ward clusters.

| Ward cluster | Prevalence of experience of dental decay (%) | Mean number of teeth with experience of dental decay in all examined children n (95% CI) | Mean number of teeth with experience of dental decay in children with any decay experience n (95% CI) |
|---------------------------|---|---|---|
| East Oldham | 59.6 | 2.6 (1.90 - 3.26) | 4.3 (3.42 - 5.22) |
| West Oldham | 56.1 | 3.0 (2.40 - 3.56) | 5.3 (4.60 - 6.02) |
| Failsworth and Hollinwood | 47.8 | 1.4 (0.50 - 2.37) | 3.0 (1.50 - 4.50) |
| Royton, Shaw and Crompton | 35.7 | 1.2 (0.42 – 2.00) | 3.4 (1.96 - 4.84) |
| Chadderton | 32.1 | 1.3 (0.72 - 1.95) | 4.2 (2.78 - 5.54) |
| Saddleworth and Lees | 18.6 | 0.7 (0.00 - 1.36) | 3.6 (0.57 - 6.68) |

Summary

In Oldham average levels of dental decay are higher than the average for England. Within Oldham, the highest levels of experience of dental decay are found in the ward clusters of East Oldham, West Oldham and also Failsworth and Hollinwood.

The small sample size means it is not possible to provide information at ward level. Future surveys could be commissioned to provide larger samples to facilitate local analysis. Commissioning High Quality Information to Support Oral Health Improvement: A toolkit about dental epidemiology for local authorities, commissioners and partners is available to support the commissioning of oral health surveys⁴.

Public health interventions can improve child oral health at a local level. Local authorities improving oral health: commissioning better oral health for children and young people is available to support local authorities to commission oral health improvement programmes for children and young people aged up to 19 years⁵.

If further local analysis is required, please contact the national dental public health team: DentalPHIntelligence@phe.gov.uk

References

- 1. Public Health England (2020). National Dental Epidemiology Programme for England: oral health survey of 5-year-olds 2019 [Online]. Available at: https://www.gov.uk/government/statistics/oral-health-survey-of-5-year-old-children-2019 [Accessed 29 May 2020].
- 2. Pine, C.M., Pitts, N.B. and Nugent, Z.J. (1997a). British Association for the Study of Community Dentistry (BASCD) guidance on the statistical aspects of training and calibration of examiners for surveys of child dental health. A BASCD co-ordinated dental epidemiology programme quality standard. Community Dental Health 14 (Supplement 1):18-29.
- 3. Children's services statistical neighbour benchmarking tool [online]. Available at: www.gov.uk/government/publications/local-authority-interactive-tool-lait [Accessed 29 May 2020].
- 4. Public Health England (2016). Commissioning high quality information to support oral health improvement. A toolkit about dental epidemiology for local authorities, commissioners and partners [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/773332/Commissioning_High_Quality_Information_to_Support_Oral_Health_Improvement.pdf
- 5. Public Health England (2014). Local authorities improving oral health: commissioning better oral health for children and young people [Online]. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/321503/CBOHMaindocumentJUNE2014.pdf

More information is available at www.gov.uk/government/collections/oral-health Please send any enquiries to DentalPHIntelligence@phe.gov.uk

PHE publications gateway number: GW-1530

© Crown Copyright, 2021. You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence