

In Medway 246 5-year-olds (approximately 77.5% of those sampled) were examined at school by trained and calibrated examiners using the national standard method².

Figure 1: Prevalence of experience of dental decay and mean number of teeth with experience of dental decay in 5-year-olds in Medway, other local authorities in the South East and England.




Table 1: Experience of dental decay in 5-year-olds in Medway, other local authorities in the South East and England.

Local authority	Prevalence of experience of dental decay (%)	Mean number of teeth with experience of dental decay in all examined children n (95% confidence intervals)	Mean number of teeth with experience of dental decay in children with any decay experience n (95% confidence intervals)
Slough	37.9	1.7 (1.31 - 2.01)	4.4 (3.74 - 5.02)
Reading	30.6	1.2 (0.92 - 1.52)	4.0 (3.35 - 4.61)
Medway	25.5	0.8 (0.54 - 1.09)	3.2 (2.44 - 3.94)
ENGLAND	23.4	0.8 (0.78 - 0.81)	3.4 (3.36 - 3.44)
Oxfordshire	21.3	0.7 (0.59 - 0.79)	3.2 (2.93 - 3.56)
Kent	19.9	0.7 (0.59 - 0.72)	3.3 (3.08 - 3.51)
Windsor and Maidenhead	19.7	0.7 (0.48 - 0.92)	3.6 (2.83 - 4.28)
Buckinghamshire	19.3	0.6 (0.49 - 0.78)	3.3 (2.77 - 3.86)
Wokingham	18.3	0.5 (0.31 - 0.66)	2.7 (2.10 - 3.25)
Milton Keynes	18.1	0.7 (0.48 - 0.95)	3.9 (3.04 - 4.85)
SOUTH EAST	17.6	0.6 (0.55 - 0.61)	3.3 (3.17 - 3.40)
West Berkshire	14.7	0.5 (0.34 - 0.74)	3.7 (2.87 - 4.46)
Hampshire	14.0	0.4 (0.36 - 0.46)	2.9 (2.70 - 3.15)
Bracknell Forest	11.5	0.5 (0.27 - 0.72)	4.3 (3.12 - 5.51)
Brighton and Hove	10.3	0.3 (0.17 - 0.41)	2.8 (2.07 - 3.62)
East Sussex	8.7	0.3 (0.21 - 0.36)	3.3 (2.71 - 3.80)

Local authority	Prevalence of experience of dental decay (%)	Mean number of teeth with experience of dental decay in all examined children n (95% confidence intervals)	Mean number of teeth with experience of dental decay in children with any decay experience n (95% confidence intervals)		
Isle of Wight	Did Not Participate				
Portsmouth	Did Not Participate				
Southampton	Did Not Participate				
Surrey	Did Not Participate				
West Sussex	Did Not Participate				

Table 2. Measures of oral health among 5-year-olds in Medway, it's statistical neighbours, i the South East and England.

	Medway	Statistical neighbour within South East: Kent	Statistical neighbour comparator 1: Swindon	South East	England
Prevalence of experience of dental decay	25.5%	19.9%	28.9%	17.6%	23.4%
Mean number of teeth with experience of dental decay	0.8	0.7	0.9	0.6	0.8
Mean number of teeth with experience of decay in those with experience of dental decay	3.2	3.3	3.2	3.3	3.4
Mean number of decayed teeth in those with experience of dental decay	2.4	2.4	2.3	2.5	2.7
Proportion with active decay	22.7%	16.4%	24.7%	14.8%	20.4%
Proportion with experience of tooth extraction ⁱⁱ	2.3%	2.2%	2.9%	1.7%	2.2%
Proportion with dental abscess	0.0%	0.1%	1.3%	0.7%	1.0%
Proportion with teeth decayed into pulp	2.2%	2.0%	4.0%	2.0%	3.3%
Proportion with decay affecting incisors ⁱⁱⁱ	3.1%	2.4%	10.6%	3.4%	5.2%
Proportion with high levels of plaque present on upper front teeth ^{iv}	0.0%	0.2%	0.7%	0.6%	1.2%


¹ generated by the children's services statistical neighbour benchmarking tool, the neighbour within the South East has "Very Close" comparator characteristics and the national neighbour 1 has "Very Close" comparator characteristics³.

ii experience of extraction of one or more teeth on one or more occasions.

iii dental decay involving one or more surfaces of upper anterior teeth.

iv indicative of poor tooth brushing habits.


Figure 2: Prevalence of experience of dental decay in 5-year-olds in Medway, by local authority Index of Multiple Deprivation (IMD) 2019 quintiles.



IMD 2019 quintiles (within local authority)


Error bars represent 95% confidence limits

Figure 3: Slope index of inequality in the prevalence of experience of dental decay in 5-yearolds in the South East.



Error bars represent 95% confidence limits

Figure 4: Prevalence of experience of dental decay in 5-year-olds in Medway, the South East and England, by year.




Error bars represent 95% confidence limits

Table 3: Experience of dental decay in 5-year-olds in the South East, by ethnic group.

Ethnic group	Number of children examined (N)	Prevalence of experience of dental decay (%)	Mean number of teeth with experience of dental decay among children with any experience of dental decay n (95% CI)	Prevalence of dental decay affecting incisors (%)
Other ethnic background	186	39.2	4.7 (3.93 - 5.52)	14.0
Asian / Asian British	845	34.3	4.0 (3.69 - 4.38)	12.7
Mixed	502	19.7	3.3 (2.74 - 3.80)	5.4
Black / Black British	249	19.7	3.9 (3.05 - 4.79)	6.0
Not provided	729	16.9	3.3 (2.85 - 3.75)	3.0
White	9,133	15.2	3.0 (2.88 - 3.14)	2.1
South East	11,644	17.6	3.3 (3.17 - 3.40)	3.4

Figure 5: Prevalence of experience of dental decay in 5-year-olds in the South East, by ethnic group.



Summary

In Medway average levels of dental decay are similar to the average for England.

The small sample size means it is not possible to provide information at ward level. Future surveys could be commissioned to provide larger samples to facilitate local analysis. Commissioning High Quality Information to Support Oral Health Improvement: A toolkit about dental epidemiology for local authorities, commissioners and partners is available to support the commissioning of oral health surveys⁴.

Public health interventions can improve child oral health at a local level. Local authorities improving oral health: commissioning better oral health for children and young people is available to support local authorities to commission oral health improvement programmes for children and young people aged up to 19 years⁵.

If further local analysis is required, please contact the national dental public health team: DentalPHIntelligence@phe.gov.uk

References

- 1. Public Health England (2020). National Dental Epidemiology Programme for England: oral health survey of 5-year-olds 2019 [Online]. Available at: https://www.gov.uk/government/statistics/oral-health-survey-of-5-year-old-children-2019 [Accessed 29 May 2020].
- 2. Pine, C.M., Pitts, N.B. and Nugent, Z.J. (1997a). British Association for the Study of Community Dentistry (BASCD) guidance on the statistical aspects of training and calibration of examiners for surveys of child dental health. A BASCD co-ordinated dental epidemiology programme quality standard. Community Dental Health 14 (Supplement 1):18-29.
- 3. Children's services statistical neighbour benchmarking tool [online]. Available at: www.gov.uk/government/publications/local-authority-interactive-tool-lait [Accessed 29 May 2020].
- 4. Public Health England (2016). Commissioning high quality information to support oral health improvement. A toolkit about dental epidemiology for local authorities, commissioners and partners [Online]. Available at:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/773332/Commissioning_High_Quality_Information_to_Support_Oral_Health_Improvement.pdf

More information is available at www.gov.uk/government/collections/oral-health Please send any enquiries to DentalPHIntelligence@phe.gov.uk

PHE publications gateway number: GW-1530

© Crown Copyright, 2021. You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence